Contents lists available at ScienceDirect

Virology

journal homepage: www.elsevier.com/locate/yviro

West African *Anopheles gambiae* mosquitoes harbor a taxonomically diverse virome including new insect-specific flaviviruses, mononegaviruses, and totiviruses



Joseph R. Fauver^a, Nathan D. Grubaugh^a, Benjamin J. Krajacich^a, James Weger-Lucarelli^a, Steven M. Lakin^a, Lawrence S. Fakoli III^b, Fatorma K. Bolay^b, Joseph W. Diclaro II^c, Kounbobr Roch Dabiré^d, Brian D. Foy^a, Doug E. Brackney^{a,1}, Gregory D. Ebel^{a,*}, Mark D. Stenglein^{a,*}

^a Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort Collins, Colorado, USA

^b Liberian Institute for Biomedical Research, Charlesville, Liberia

^c United States Naval Medical Research Unit No. 3, Cairo, Egypt

^d Institut de Recherche en Sciences de la Santé, Bobo-Dioulasso, Burkina Faso

ARTICLE INFO

SEVIEI

Article history: Received 4 May 2016 Returned to author for revisions 29 July 2016 Accepted 31 July 2016 Available online 15 September 2016 Keywords: Anopheles gambiae Arthropod viruses Metagenomics Surveillance Bioinformatics Flavivirus Mononegavirus Totivirus

ABSTRACT

Anopheles gambiae are a major vector of malaria in sub-Saharan Africa. Viruses that naturally infect these mosquitoes may impact their physiology and ability to transmit pathogens. We therefore used metagenomics sequencing to search for viruses in adult *Anopheles* mosquitoes collected from Liberia, Senegal, and Burkina Faso. We identified a number of virus and virus-like sequences from mosquito midgut contents, including 14 coding-complete genome segments and 26 partial sequences. The coding-complete sequences define new viruses in the order Mononegavirales, and the families Flaviviridae, and Totiviridae. The identification of a flavivirus infecting *Anopheles* mosquitoes broadens our understanding of the evolution and host range of this virus family. This study increases our understanding of virus diversity in general, begins to define the virome of a medically important vector in its natural setting, and lays groundwork for future studies examining the potential impact of these viruses on anopheles biology and disease transmission.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Virus discovery Pathogen discovery Virome

Mosquitoes (Diptera: Culicidae) are the most important vectors of human disease. *Anopheles gambiae* and other anopheline mosquitoes are the vectors of *Plasmodium* parasites in Africa, which cause nearly 200 million malaria cases annually on the continent (World Health Organization, 2015). These mosquitoes are also responsible for transmitting O'nyong-nyong virus, a human pathogenic alphavirus capable of causing large disease outbreaks

* Corresponding authors.

E-mail addresses: Gregory.Ebel@colostate.edu (G.D. Ebel),

Mark.Stenglein@colostate.edu (M.D. Stenglein).

(Lanciotti et al., 1998; Lutwama et al., 1999). In addition, *Anopheles* spp. mosquitoes are vectors of *Wuchereria bancrofti* roundworms, the causative agent of lymphatic filariasis, which affects over 100 million people in sub-Saharan Africa (World Health Organization, 2016). Current interventions for these diseases are inadequate, and future strategies need to deploy current and novel interventions to disrupt pathogen transmission.

Constituents of arthropod microbiomes are being increasingly scrutinized for their potential to alter the arthropod's ability to transmit co-infecting human pathogens (Weiss and Aksoy, 2011). The microbiome of *Anopheles* species, especially in the gut, has been shown to be quite diverse and can vary depending on the mosquito's environment (Buck et al., 2016; Grubaugh et al., 2015). Furthermore, certain bacteria can influence *Plasmodium* development in the mosquito (Boissière et al., 2012; Cirimotich et al., 2011;

0042-6822/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

¹ Present address: Center for Vector Biology and Zoonotic Diseases, Connecticut Agricultural Experiment Station, New Haven, Connecticut, USA.

Dong et al., 2009; Gendrin et al., 2015; Hughes et al., 2014). During the last decade there has been increased attention to insect-specific viruses (ISVs) and their potential role in disrupting pathogen transmission (Blitvich and Firth, 2015; Bolling et al., 2015; Junglen and Drosten, 2013; Vasilakis and Tesh, 2015). The majority of ISVs have been described in mosquitoes, although they are known to occur in several arthropod orders, including Hemiptera (i.e. true bugs) (Li et al., 2015) and Parasitiformes (e.g. ticks) (Tokarz et al., 2014). ISVs belong to taxonomically diverse virus families including Bunyaviridae (Auguste et al., 2014; Chandler et al., 2014; Marklewitz et al., 2011; Marklewitz et al., 2013; Yamao et al., 2009), Flaviviridae (Blitvich and Firth, 2015; Cook et al., 2006; Hoshino et al., 2007; Lee et al., 2013; Misencik et al., 2016; Tyler et al., 2011), Reoviridae (Attoui et al., 2005; Auguste et al., 2015; Hermanns et al., 2014) Rhabdoviridae (Kuwata et al., 2011; Quan et al., 2010; Vasilakis et al., 2014a), and Togaviridae (Nasar et al., 2012). In addition, ISVs in the Birnaviridae (Huang et al., 2013; Marklewitz et al., 2012; Vancini et al., 2012), Nodaviridae (Schuster et al., 2014), Tymoviridae (Wang et al., 2012), and Parvoviridae (Ren



Fig. 1. Mosquito collection sites in Liberia, Senegal, and Burkina Faso. In Liberia, mosquitos were sampled from 6 villages within an area with an approximate radius of 16 km. The collection site in Burkina Faso for mosquito eggs that were used to found a laboratory colony is indicated.

Table 1

Summary of mosquito samples analyzed.

et al., 2008) families have been characterized, the later from which an *Anopheles*-specific densovirus is being examined as a paratransgenesis candidate (Ren et al., 2008; Suzuki et al., 2014). Recently discovered ISVs include those in the family *Mesoniviridae* (Kuwata et al., 2013; Nga et al., 2011; Vasilakis et al., 2014b; Warrilow et al., 2014; Zirkel et al., 2011; Zirkel et al., 2013) and a variety of positive sense ssRNA viruses including the negeviruses (Kallies et al., 2014; Nabeshima et al., 2014; Vasilakis et al., 2013). Recently, two new RNA viruses, a dicistrovirus and a cypovirus, were identified in *Anopheles* species mosquitoes (Carissimo et al., 2016). Mosquitoes are divided into two subfamilies: *Culicinae* and *Anophelinae*, however, a disproportionate amount of mosquito ISVs have been identified from culicine mosquitoes, leaving anopheline mosquitoes relatively understudied.

We therefore used metagenomic sequencing to identify viruses infecting wild *An. gambiae*, *An. funestus*, and *An. rufipes* mosquitoes in West Africa. We sampled adult mosquitoes from villages in rural Burkina Faso, Liberia, and Senegal, sequenced RNA, and searched datasets for virus sequences. We identified sequences from multiple new viruses. For several of these, we generated coding complete genome sequences, performed comparative and phylogenetic analyses, and determined the prevalence from our fieldcollected samples. Our findings indicate that anopheline mosquitoes naturally harbor multiple viruses including flaviviruses.

2. Materials and methods

Mosquito samples used in this study were collected from 2012 to 2015 on separate trips to Senegal, Liberia, and Burkina Faso (Fig. 1 and Table 1). Indoor resting bloodfed mosquitoes were collected in Senegal (Alout et al., 2014; Krajacich et al., 2015) and Liberia (Grubaugh et al., 2015) as previously described. Mosquitoes from Burkina Faso were colonized and reared at the Colorado State University (CSU) Arthropod-Borne and Infectious Disease Laboratory prior to being sampled.

2.1. Sample preparation

2.1.1. Burkina Faso

A laboratory colony of *An. gambiae s.s.* was established by the Institut de Recherche en Sciences de la Santé from larvae collected

Sample set	Location	Date	Number mosquitos analyzed	Mosquito species ^a	Analysis method		
1	Senegal	8/22/2012	41 ^b	An. funestus/An. gambiae/An. rufipes	NGS (MiSeq)		
2	Burkina Faso	1/5/2015 ^c	17	An. gambiae	NGS (NextSeq)		
3	66	12/30/2015 ^c	3	"	"		
4	Liberia (village A)	6/11&13/2013	31	An. gambiae	NGS (HiSeq)		
5	66	6/15/2013	38	ц. ц	66		
6	"	6/19/2013	24	"	"		
7	"	6/21/2013	41	"	"		
8	66	6/23/2013	29	"			
9	"	6/25/2013	57	и	66		
10	Liberia (village B)	6/10&14/2013	51	и	66		
11	"	6/22&26/2013	57	и	66		
12	Liberia (village C)	3/30/2015	15 ^d	An. gambiae	PCR / Sanger sequencing		
13	Liberia (village D)	3/31/2015	15	"	"		
14	Liberia (village E)	4/1/2015	15	и	66		
15	Liberia (village A)	4/2/2015	15	и	66		
16	Liberia (village F)	4/3/2015	15	"	"		

Notes:

^a As determined by field identification and molecular analysis (see Section 2).

^b Mosquitos in sample sets 1-11 were pooled (1 pool per sample set) for sequencing.

^c These mosquitoes were sampled on these dates from a laboratory colony that was derived from *An. gambiae* larvae collected in Burkina Faso.

^d Mosquitos in sample sets 12–16 were analyzed individually (i.e. not pooled) using PCR and Sanger sequencing to validate NGS results and measure prevalence.

in Burkina Faso in 2014 and eggs from this colony were subsequently shipped to Colorado State University (CSU). Mixed sex, non-bloodfed mosquitoes from the colony at CSU were homogenized in 1 ml of mosquito diluent (80% PBS, 20% FBS, supplemented with penicillin, streptomycin, gentamicin, and amphotericin B (Fauver et al., 2015)) with a steel ball bearing for RNA extractions. 50 μ l of cleared supernatant was used for RNA extraction with the Mag-Bind Viral DNA/RNA kit (Omega, Georgia, USA) with the KingFisher Flex Magnetic Particle Processor (Thermo Fisher Scientific, Massachusetts, USA) according to manufacturer's protocol. Libraries were prepared using the Ovation RNA-Seq System V2 (NuGEN, California, USA) and Ovation Ultralow DR Multiplex System 1–96 (NuGEN) and sequenced on an Illumina NextSeq at the CSU NGS facility.

2.1.2. Senegal

Field-caught bloodfed mosquitoes were pooled by date and stored in RNA Later (Ambion) at -80 °C. Pools were thawed and RNA Later was removed. 1 ml of PBS was added and mosquito pools were centrifuged for 5 min. 140 µl of supernatant was used for RNA extraction using Qiagen Viral RNA Mini Kit (Qiagen, Hilden, Germany) according to manufacturer's protocol with slight modification. Extracted RNA was subjected to DNase treatment (Thermo Fisher Scientific) and purification using Agencourt RNAclean XP beads (Bechman Colter Genomics, Pasadena, CA). cDNA was amplified using the Ovation RNA-Seq System V2 and prepared for library construction using the Ovation Ultralow DR Multiplex System 1–96 as described (Grubaugh et al., 2016). Libraries were sequenced on a NextSeq instrument at the CSU NGS facility.

2.1.3. Liberia

Mosquitoes were collected and processed as previously described (Grubaugh et al., 2015). Briefly, bloodfed mosquitoes were knocked down with triethylamine and bloodmeals were expelled onto Whatman Flinders Technology Associates (FTA) clone saver cards (GE Healthcare Life Sciences, Little Chalfont, United Kingdom). Cards were stored at -20 °C and shipped to CSU for additional processing. Mosquito dried blood spots (M-DBS) were removed using a Harris 3 mm micro-puncher (GE Healthcare Life Sciences) and placed into RNA Rapid Extraction Solution (Ambion, Texas, USA) to elute nucleic acid off of the cards. RNA was extracted using the Mag-Bind Viral DNA/RNA kit with the KingFisher Flex Magnetic Particle Processor according to manufacturer's protocol. Libraries were prepared using the Ovation RNA-Seq System V2 and Nextera XT DNA Sample Preparation Kit (Illuminia, San Diego, CA) and sequenced on an Illumina HiSeq (Beckman Colter Genomics, Danvers, MA).

2.2. Sequence analysis

Sequencing datasets were processed with the goal of taxonomically assigning non-mosquito reads. First, low quality and Illumina adapter sequences were removed using the Trimmomatic tool version 0.32 with the following settings: ILLUMINACLIP: NexteraPE:1:30:10:4: true LEADING:20 TRAILING:20 SLI-DINGWINDOW:4:25 MINLEN:60 (Bolger et al., 2014). Potential PCR duplicate sequences were collapsed using the CD-HIT-EST tool, version 4.6 with parameter -c 0.96 (Li and Godzik, 2006). Then, mosquito sequences were removed by aligning reads to databases of An. gambiae genomic sequences using Bowtie2 version 2.2.5, with parameters –sensitive –score-min C,60,0 (Holt et al., 2002). Remaining sequences were de novo assembled into contiguous sequences using the SPAdes genome assembler (Bankevich et al., 2012; Langmead and Salzberg, 2012). Resulting contigs and nonassembling reads were then taxonomically assessed, first by using the gnapl tool, version 2014-12-28, to align to the NCBI nt nucleotide database (Wu and Nacu, 2010). Sequences that did not produce a nucleotide-level alignment were then searched via translated-nucleotide to protein alignments against the NCBI nr protein sequence database using the Rapsearch2 tool, version 2.23, with parameters –a t, – 1 20, and –e1 e-2 (Zhao et al., 2012). Draft virus genome sequences were validated by mapping individual reads to assemblies using Bowtie2 as above and in some cases using PCR and Sanger sequencing. Resulting alignments were imported into Geneious software version 9.0.4 and manually inspected (Kearse et al., 2012). Sequence datasets have been deposited in the NCBI Short Read Archive (SRA) with accession PRJNA327220.

2.3. Analysis of predicted viral protein sequences

ORFs in viral genome assemblies were predicted using Geneious software. Homologs of predicted protein sequences were detected using the BLASTP tool (version 2.2.25+) to search the NCBI non-redundant protein database (nr) (Altschul et al., 1997). For sequences with no detectable similarity by BLASTP, the HHpred homology detection and structure prediction tool (version 2.0) was also used (Soding et al., 2005). The transmembrane prediction tool in Geneious was used to predict transmembrane domains. Virus and virus-like contigs longer than 500 nt long were aligned to the NCBI nr protein sequence database using BLASTX to determine taxonomic classification, closest relative, and percent similarity to closest related sequences.

2.4. Phylogenetic analysis

Predicted viral protein sequences were used to query the NCBI nr protein database using the BLASTP tool (Altschul et al., 1990). Database sequences that aligned with an E-value less than 10^{-3} and that were full length or nearly full-length were downloaded. These were collapsed using the CD-HIT tool, version 4.6 using parameter -c 0.9 (Li and Godzik, 2006). These representative sets of sequences were aligned using the MAFFT software, version 7.221, using the LINSI mode (Katoh and Standley, 2013). Phylogenetically uninformative columns were removed from multiple alignments using the GBlocks tool, version 0.91b, with parameter b5=n (Talavera and Castresana, 2007). These trimmed alignments were used to create phylogenies with the MrBayes version 3.2.5 with commands preset aamodelpr=mixed and mcmc ngen= 1,000,000 (Ronquist and Huelsenbeck, 2003). Convergence was confirmed by inspecting the standard deviation of split frequencies. Phylogenies were visualized using FigTree version 1.4.2 (http://tree.bio.ed.ac.uk/software/figtree/).

2.5. Validation of mosquito species collected

We used a molecular strategy to corroborate field identification of the species of collected mosquitoes. We collected a representative set of mosquito cytochrome c oxidase subunit 1 (CO1) gene sequences (Supplemental Table 1) and used Bowtie2 to align sequences from our quality-filtered datasets to these sequences and tabulated the fractions of reads aligning to each mosquito species. This analysis corroborated field-based species identifications (Table 1).

2.6. Prevalence of Anopheles viruses

The abundance of virus reads in sequencing datasets was calculated by using Bowtie2 to map all unique, host-filtered reads to all of the virus sequences identified in this study. The abundance was defined as the number of mapping reads per million unique reads in each dataset.

Field prevalence of Anopheles Flavivirus, Bolahun Anopheles virus, and Anopheles Totivirus was determined using independent sample sets collected in 2015 (Table 1). RNA isolated from individual Liberian M-DBS was tested for the presence of each virus by RT-PCR using the Qiagen OneStep RT-PCR kit according to manufacturer's protocol with virus specific primers (Supplemental Table 2). 15 M-DBS from 5 villages (75 total) were used to calculate field prevalence.

3. Results

Α

We collected adult Anopheles mosquitoes from several West Africa locations and performed metagenomic sequencing of midgut RNA-derived libraries to search for viruses. In total, we analyzed 328 adult An. gambiae from Liberia and 41 mixed An. gam*biae/An. funestus/An. rufipes* from Senegal ($\sim 1/3$ of each species). These mosquitoes were collected over the course of several years from multiple villages (Table 1). We also analyzed 20 mosquitoes from a laboratory colony established in 2014 from Burkina Faso An. gambiae s.s. larvae. These colony mosquitoes were sampled in 2 batches \sim 12 months apart. Mosquitoes were combined into pools of between 3 and 57 mosquitoes each, and pools were sequenced on Illumina instruments to a median depth of 4.8×10^7 150 nt read pairs per pool. After filtering low quality, duplicate, and mosquito-derived reads, a median of 3.0×10^5 reads remained (0.6%). Remaining reads were de novo assembled and taxonomically assigned by comparison to sequences in Genbank, first by nucleotide-to-nucleotide alignments, then by translated-nucleotide to protein alignments. We identified a number of putative viral sequences and determined coding-complete genome segment sequences where possible (Ladner et al., 2014). Sequencing depth was determined for each coding complete genome (Supplemental Fig. 1). We also used sequencing data to confirm the species composition of the collected mosquito sample sets (Table 1).

3.1. Anopheles flavivirus

We identified flavivirus sequences in datasets from mosquitoes collected in Liberia and Senegal (Fig. 2 and Table 1). The genus Flavivirus (family Flaviviridae) includes viruses whose life cycle involves alternating replication in vertebrate and arthropod hosts and members whose life cycle is restricted to one host or the other (Van Regenmortel, 2000a). Typical flaviviruses have positive-sense ssRNA genomes that encode a single large polyprotein. We



Fig. 2. Anopheles flavivirus genome organization, dinucleotide usage, and phylogeny (A) Genomic organization of anopheles flavivirus. Predicted functional domains of the viral polyprotein are indicated as is the fifo ORF and the predicted ribosomal frameshift "slippery" sequence at its beginning is indicated by a triangle. The other AnFV and AnFV-like sequences identified in this study (KX148547-KX148549) have a similar organization. (B) AnFV clusters phylogenetically with classic ISFVs. A multiple sequence alignment of flavivirus NS5 protein sequences was used to create a Bayesian phylogeny. The phylogeny was rooted on the branch to Tamana bat virus (not shown). Posterior probabilities of select nodes are indicated. A fully-labeled version of this phylogeny including accession numbers and node posterior probabilities is available as Supplemental Fig. 3. (C) Phylogeny as in (B) but focused on the clSFV clade. (D) Dinucleotide usage in AnFV supports its categorization as a classic insect specific flavivirus (ISFV). CpG and UpA dinucleotide frequencies for viruses in the genus Flavivirus in the NCBI RefSeq database are indicated. Points are color coded according to flavivirus categories as indicated and as in (Blitvich and Firth, 2015). FV: flavivirus. Anopheles flavivirus points are indicated.

assembled two coding-complete flavivirus genomes from Liberia datasets, which shared 95% pairwise nucleotide identity. We designated these as Anopheles flavivirus (AnFV) – variants 1 and 2. We also assembled partial sequences from at least one additional flavivirus from the Senegal datasets, which we called Anopheles flavivirus-like sequences 1 and 2 (Table 1). Anopheles flavivirus-like sequences 1 shared ~79% pairwise nucleotide identity with the AnFV sequences. These and other genome sequences represent the consensus of sequences that shared \geq 99.5% pairwise identity. Single nucleotide variants were evident in the various datasets (Supplemental Fig. 2). Given that these datasets derive from pools of mosquitoes, this variation could represent intra- or inter-host diversity, or both.

The genome organization, gene content, phylogenetic placement, and dinucleotide usage supports the classification of AnFV as a "classic" insect-specific flavivirus (cISFV) (Blitvich and Firth, **2015**; **Bolling et al.**, 2015). The AnFV genome contains an ORF of 10032 nt predicted to encode a polyprotein of 3341 amino acids (Fig. 2A). The polyprotein is predicted to be co- and post-transcriptionally cleaved to produce the typical 3 structural and 7 non-structural flavivirus proteins and we identified putative cleavage sites. The polyprotein shares 35–43% global pairwise amino acid identity with sequences from other cISFVs.

In addition to the polyprotein ORF, the genome contains an 840 nt ORF overlapping with the NS2 coding region (Fig. 2A). The reading frame of this ORF is -1 relative to the polyprotein ORF, and 5 nt downstream of the predicted NS2A cleavage site, a putative "slippery" sequence (GGAUUUU) was identified as a likely site of ribosomal frameshifting. These "fifo" ORFs (fairly interesting flavivirus ORF) are a characteristic of clSFV genomes (Blitvich and Firth, 2015; Firth et al., 2010). The predicted AnFV FIFO protein possesses no detectable sequence similarity with other clSFV FIFO



Fig. 3. Bolahun and Gambie viruses. (A) Genomic organization of Bolahun virus. Predicted coding sequences are indicated. Gambie virus has essentially identical genome organization. (B) Bolahun and Gambie viruses cluster with bornaviruses, nyamiviruses, and arthropod viruses. RdRp-based Bayesian phylogeny. Posterior probabilities of select nodes are indicated. Phylogenies showing all mononegavirus sequences are shown in Supplemental Figs. 4 and 5. (C) Viruses in this virus clade encode a predicted small zinc finger protein just upstream of the L ORF. Genome cartoons of Bolahun virus, Xincheng mosquito virus, and Shuangao fly virus 2. Open reading frames are labeled according to (Li et al., 2015). ORFs not annotated in KM817661 and KM817638 are outlined with dotted lines. Small ORFs predicted to encode Zn finger proteins upstream of L ORFs are colored purple. (D) Multiple alignment of proteins encoded by small ORFs highlighted in (C). Cysteine CXXC motifs characteristic of Zn finger domains are underlined.



proteins. However, like other cISFV FIFO proteins, the AnFV protein contains predicted transmembrane domains (Firth et al., 2010). Phylogenetic analysis corroborated the designation of AnFV as a cISFV (Fig. 2**B**, **C** and Supplemental Fig. 3). In Bayesian phylogenies based on alignments of NS5 protein sequences, the AnFV sequences occupy a well-supported branch within the cISFV clade. Analyses based on alignments of full polyprotein sequences produced phylogenies with essentially identical topologies.

Viral genomes often exhibit patterns of dinucleotide usage similar to that of their hosts. On this basis, cISFV genomes can be distinguished from those of flaviviruses whose lifecycles include replication in vertebrates (Blitvich and Firth, 2015). Based on dinucleotide usage, AnFV clusters with cISFVs (Fig. 2**D**).

3.2. Bolahun and Gambie viruses

We identified mononegavirus sequences in mosquitoes from Liberia and Senegal and from our Burkina Faso-derived colony. *Mononegavirales* is a large and diverse order of viruses that have single-stranded negative polarity RNA genomes and evolutionarily related RNA dependent RNA polymerase (RdRp) genes (Van Regenmortel, 2000b). We assembled apparently coding-complete mononegavirus genomes from each of these three datasets (Fig. 3). The genomes from Liberia and Burkina Faso were closely related (94% pairwise nt identity), and we designated these as Bolahun virus (BOAV) – variants 1 and 2. The sequence from Senegal, which we named Gambie virus (GAMV), shared ~60% pairwise nt identity with BOAV.

The BOAV and GAMV genomes share a similar overall genome organization, with 6 non-overlapping ORFs (Fig. 3A). In mononegaviruses, the ORF nearest the 3' end of the genome encodes the viral nucleoprotein. The BOAV and GAMV ORF1 is predicted to encode protein of 446 amino acids (lengths given for BOAV, accession KX148552) with no transmembrane domains and an isoelectric point of 8.7. By BLASTP search, the only identifiable homologous sequence was that encoded by Xincheng mosquito virus ORF1 (Li et al., 2015), which shares 19% pairwise global amino acid identity. These are likely nucleoproteins for these viruses. ORF2 encodes a predicted small transmembrane domaincontaining protein of 65 residues. ORF3 encodes a protein predicted to be 446 AA long with no detectable similarity to known proteins by BLASTP or HHPRED analyses w/ E-value cutoff 0.1 (Soding et al., 2005). ORF4 encodes the predicted viral glycoprotein, a 638 AA protein with 3 transmembrane domains and sequence similarity to a variety of mononegavirus glycoproteins. As with the putative nucleoprotein, the most similar sequence is from Xincheng mosquito virus, with 38% global pairwise identity. As in all mononegaviruses, the last ORF is predicted to encode the large RdRp (L) protein. BOAV and GAMV proteins share between 36% (ORF2 protein) and 72% (glycoprotein) global amino acid identity.

In phylogenies based on alignments of L protein sequences, BOAV and GAMV form a well-supported clade with Xincheng mosquito virus and Shuangao fly virus, which were identified in samples from China from an *Anopheles sinensis* mosquito and a *Psychoda alternata* fly (Fig. 3**B and** Supplemental Figs. 4 and 5). These viruses form a sister clade with those of the *Bornaviridae* and the *Nyamiviridae*, two mononegavirus families (Kuhn et al., 2013; Kuhn et al., 2015; Mihindukulasuriya et al., 2009).

In the 3 BOAV and GAMV genomes, and in the genomes of Xincheng mosquito virus and Shuangao fly virus 2, there is a small ORF upstream of the L ORF (Fig. 3C). These ORFS range from 123 to 162 nt and encode predicted proteins of 40–53 AA with CXXC motifs characteristic of zinc ribbon type zinc finger domains (Krishna et al., 2003) (Fig. 3D). By HHPRED analysis, similarity to various cellular zinc finger domains was detected. In addition to their phylogenetic placement, a defining characteristic of the

viruses in this clade may be the presence of these small ORFs.

3.3. Anopheles totivirus

We identified a totivirus-like sequence in An. gambiae mosquitoes in Liberia (Fig. 4A). Totiviruses were historically only known to infect plant and protist hosts. However a growing number of reports have described totiviruses infecting or associated with arthropod hosts (Dantas et al., 2016; Isawa et al., 2011; Koyama et al., 2015; Martinez et al., 2016; Poulos et al., 2006; Wu et al., 2010; Zhai et al., 2010). Totiviruses have dsRNA genomes that typically have 2 overlapping ORFs that encode the viral capsid and RdRp proteins. We assembled the apparent coding-complete genome of this virus, which we designated Anopheles totivirus (AToV). There are two large ORFs in the AToV genome, but they are in the same reading frame and their translation is unlikely to involve ribosomal frameshifting (Fig. 4A). The first ORF encodes a predicted protein of 980 AA with similarity to sequences encoded in several dipteran genomes. The second ORF encodes the predicted viral RdRp of 1006 AA. In phylogenies based on alignments of totivirus RdRp sequences, AToV clusters with arthropod-infecting or arthropodassociated totiviruses (Fig. 4B).

3.4. Other virus and virus-like sequences

In addition, we identified a number of other putative virusderived contiguous sequences (contigs) in our datasets (Table 2 and Fig. 5). Some of these are partial sequences and some appear to be coding complete. We restricted our analysis to contigs longer than 500 nt. In addition to the BOAV and GAMV sequences, we identified 5 additional partial mononegavirus sequences from one or more viruses present in various Liberian mosquito pools. Like BOAV and GAMV, the closest related sequences were to various mononegaviruses from Chinese mosquitoes (Table 2) (Li et al., 2015). We also identified 6 sequences most similar to various positive-sense ssRNA viruses identified in wild-caught drosophila. In addition, we identified a number of sequences with similarity to various dsRNA viruses, including partitiviruses, chrysoviruses, and endornaviruses. These sequences were generally closest to sequences from dsRNA viruses identified in arthropods (Chandler et al., 2015; Cook et al., 2013; Spear et al., 2010). The lower coverage and partial nature of these other viral sequences suggests that they might be less prevalent in the mosquito populations that we sampled, that viral nucleic acid is present at lower levels in mosquito midguts, or that sample processing and library preparation protocols may have reduced their representation in sequencing libraries (Fig. 5).

We also identified virus sequences that likely originated from laboratory contamination. These included West Nile virus (WNV) and Phasi Charoen virus (PCV) sequences, which were nearly identical to database sequences. Our laboratories routinely work with WNV, and PCV has been found in sequencing data from mosquito cell lines in our lab. Two sequencing pools contained reads to PCV sequences, and these were the only pools that had CO1 reads aligning to mosquito genera other than *Anopheles*, corroborating the designation of these sequences as laboratory contaminants.

3.5. Prevalence of viruses in wild mosquito populations

We performed RT-PCR on an independent set of samples to validate our NGS finding and to measure the prevalence of the viruses we found in Liberian mosquitoes, namely AnFV, AToV, and BOAV (Table 3 and Fig. 5). These samples were collected from additional Liberian villages ~ 2 years after the initial sample sets



Fig. 4. (A) Genomic organization of anopheles totivirus. The two large ORFs are indicated. Although the genome is depicted in the positive sense, it is presumed to be dsRNA. (B) Anopheles totivirus clusters with viruses identified in other arthropods. A multiple sequence alignment of totivirus RdRp protein sequences and related sequences was used to create a Bayesian phylogeny as described in Section 2. Phylogeny is arbitrarily rooted. Posterior probabilities of select nodes are indicated. Sequences associated with arthropods are colored purple. Clades corresponding to ICTV-classified genera are indicated.

were collected (Table 1). In the villages, prevalence for AnFV and AToV varied between 0–20% and 0–6.7%, respectively. In total, AnFV RNA was detected in 12% of sampled mosquitoes and AToV

RNA in 1.3%. BOAV, which was detected in fewer NGS datasets (Fig. 5), was not detected in any individual M-DBS from this independent set of samples.

Table 2

summary of identified virus sequences.

Name ^a	Closest Related Sequence ^b	Seq. Length (nt)	Accession	% Identity ^c	Preliminary classification
Anopheles flavivirus – variant 1	Palm Creek Virus (AGG76014.1)	10,604	KX148546	40	Flavivirus
Anopheles flavivirus – variant 2	Palm Creek Virus (AGG76014.1)	10,529	KX148547	40	Flavivirus
Anopheles flavivirus-like sequence	Nienokoue virus (YP_009041466.1)	6063	KX148548	48	
Anopheles flavivirus-like sequence	Parramatta River virus (YP_009164029.1)	2730	KX148549	34	
Anopheles totivirus	Camponotus yamaokai virus (YP_009143313.1)	6364	KX148550	30	Totivirdae
Bolahun virus – variant 1	Xincheng Mosquito virus (AJG39227.1)	11,666	KX148551	39	Mononegavirales
Bolahun virus – variant 2	Xincheng Mosquito virus (AJG39227.1)	11,672	KX148552	39	Mononegavirales
Gambie virus	Xincheng Mosquito virus (AJG39227.1)	11,652	KX148553	38	Mononegavirales
Chaq virus-like sequence 1	Chaq virus (AKH40308.1)	1361	KX148554	38	
Chaq virus-like sequence 2	Chaq virus (AKH40308.1)	1342	KX148555	32	
Chaq virus-like sequence 3	Chaq virus (AKH40308.1)	1342	KX148556	33	
Partitivirus-like sequence 1	Penicillium aurantiogriseum partiti-like virus (YP_009182157.1)	1418	KX148575	37	
Partitivirus-like sequence 2	Penicillium aurantiogriseum partiti-like virus (YP 0091821571)	1705	KX148576	38	
Partitivirus-like sequence 3	Penicilium aurantiogriseum partiti-like virus	1705	KX148577	38	
Partitivirus-like sequence 4	Heterobasidion partitivirus 8 (AFW178111)	1955	KX148578	35	
Partitivirus-like sequence 5	Grapevine partitivirus (AFX730191)	2078	KX148579	49	
Chrysovirus-like sequence	Anthurium mosaic-associated virus (ACU115631)	949	KX148557	39	
ssRNA virus-like sequence 1	Muthill virus (AM0032231)	3065	KX148580	32	
ssRNA virus-like sequence 2	Muthill virus (AM0032231)	1974	KX148581	38	
ssRNA virus-like sequence 3	Muthill virus (AMO03223.1)	1039	KX148582	29	
ssRNA virus-like sequence 4	Muthill virus (AM003223.1)	634	KX148583	34	
ssRNA virus-like sequence 5	Blackford virus (AMO03220.1)	4642	KX148584	45	
ssRNA virus-like sequence 6	Buckhurst virus (AMO03221.1)	7413	KX148585	28	
dsRNA virus-like sequence 1	dsRNA virus environmental sample (AIT39583.1)	1886	KX148558	41	
dsRNA virus-like sequence 2	dsRNA virus environmental sample (AIT39584.1)	1434	KX148559	32	
dsRNA virus-like sequence 3	dsRNA virus environmental sample (AIT39583.1)	1230	KX148560	56	
dsRNA virus-like sequence 4	dsRNA virus environmental sample (AIT39584.1)	809	KX148561	44	
dsRNA virus-like sequence 5	dsRNA virus environmental sample (AIT39584.1)	735	KX148562	30	
dsRNA virus-like sequence 6	uncultured RNA virus (AGW51771.1)	664	KX148563	28	
dsRNA virus-like sequence 7	dsRNA virus environmental sample (AJT39583.1)	634	KX148564	62	
dsRNA virus-like sequence 8	dsRNA virus environmental sample (AJT39583.1)	596	KX148565	38	
dsRNA virus-like sequence 9	dsRNA virus environmental sample (AJT39589.1)	588	KX148566	41	
Endornavirus-like sequence 1	Phaseolus vulgairs endornavirus 2 (BAM68540.1)	2858	KX148567	22	
Endornavirus-like sequence 2	Rhizoctonia cerealis endornavirus 1 (YP_008719905.1)	672	KX148568	39	
Endornavirus-like sequence 3	Bell pepper endornavirus (YP_004765011.1)	514	KX148569	39	
Mononegavirus-like sequence 1	Xincheng Mosquito virus (AJG39226.1)	964	KX148570	70	
Mononegavirus-like sequence 2	Xincheng Mosquito virus (AJG39227.1)	948	KX148571	34	
Mononegavirus-like sequence 3	Wuhan Mosquito virus 9 (AJG39215.1)	878	KX148572	56	
Mononegavirus-like sequence 4	Wuhan Mosquito virus 2 (AJG39297.1)	869	KX148573	39	
Mononegavirus-like sequence 5	Wuhan Mosquito virus 2 (AJG39331.1)	550	KX148574	45	

Notes:

^a Virus and virus-like contigs with length > = 500 nt are shown.

^b Closest relative determined by highest scoring alignment from BlastX search of NCBI nr database.

^c Pairwise percent amino acid identity of BlastX alignment to closest relative.

4. Discussion

In this study we sampled anopheline mosquitoes from Liberia and Senegal and a laboratory colony derived from Burkina Faso. We identified a number of previously undescribed virus and viruslike sequences. For some of these, we detected the same or related virus genotypes in mosquitoes that were collected in multiple countries and in mosquitoes that were sampled up to 2 years apart. Thus, infection by these viruses appears to be geographically widespread and temporally stable in mosquito populations.

It is likely that many of these sequences derive from viruses that genuinely infect these mosquitoes, although this remains to be proven. However, mosquitoes have a diverse microbiome and it cannot be ruled out that some of these sequences are derived from viruses infecting commensal organisms. And, because both male and female mosquitoes feed on plant nectar, and females take blood meals, it is also possible that some of these sequences have a dietary or environmental origin. These sequences do not appear to derive from integrated endogenous viral elements, given their absence in the *An. gambaie* genome, non-uniform presence in mosquitoes, and intact ORFs (Crochu et al., 2004). Several pieces of evidence point to AnFV, BOAV, and GAMV as legitimate mosquito viruses, including their phylogenetic placement and genomic characteristics. While groups of flaviviruses and mononegaviruses are accepted as insect-infecting viruses, totiviruses were classically thought to infect plants and fungi. AToV groups phylogenetically with totiviruses described from various arthropods, however we cannot rule out the possibility that AToV infects commensal fungi or protists of *Anopheles* mosquitoes. In any case, AnFV, BOAV, GAMV, and AToV represent legitimate viral constituents of the microbiome of *Anopheles* species mosquitoes.

Although we identified several virus sequences that likely derive from laboratory contamination, several lines of evidence support the conclusion that the other virus and virus-like sequences we identified were not contaminants. First, we observed sequence variability: for example, the AnFV sequences from the



Fig. 5. Abundance of virus and virus-like sequences in sample sets. The relative abundance of virus-mapping reads in each NGS dataset is shown (see Table 2). Virusmapping reads per million unique filtered reads are shown. Genbank accessions of virus or virus-like sequences are indicated.

Table 3					
Prevalence of	of viruses	in	Liberian	mosquitoes.	

.....

Village C		Villa	Village D		Villa	Village E		Village A			Village F			Total		
Virus AnFV BOAV AToV	+ ^a 2 0 1	N ^b 15 15 15	% pos 13.3 0 6.7	+ 3 0 0	N 15 15 15	% pos 20 0 0	+ 2 0 0	N 15 15 15	% pos 13.3 0 0	+ 0 0 0	N 15 15 15	% pos 0 0 0	+ 2 0 0	N 15 15 15	% pos 13.3 0 0	12% 0% 1.3%

^a +: number positive blood spots by RT-PCR.

^b N: number individual mosquito blood spots tested.

8 Liberian datasets, which were 99.5–100% identical to the consensus sequence, had distinct sets of single nucleotide variants (KX148546, Supplemental Fig. 2). Second, sequences had variable abundances and non-uniform presence in our datasets (Fig. 5). Third, we did not detect these sequences in other NGS datasets generated in our labs, nor were they present in online sequence databases.

The AnFV sequences we identified represent the first description of flaviviruses from anopheline mosquitoes. With the exception of tick-borne flaviviruses and no known vector flaviviruses, these viruses have been described exclusively from culicine mosquitoes (Kuno et al., 1998). The sequence of AnFV is highly divergent from other known flaviviruses, including from other clSFVs (Fig. 2**C** and Supplemental Fig. 3). This is not unexpected, as clSFVs tend to group by host mosquito genera (Blitvich and Firth, 2015). The description of AnFV further exhibits the diversity of the genus *Flavivirus* and supports the hypothesis that flaviviruses are under sampled (Cook et al., 2012; Pybus et al., 2002). It remains to be determined how AnFV will contribute to the understanding of the evolution of clSFVs, especially in regards to co-divergence (Cook et al., 2012). In any case, the additional genomic and phylogenetic information provided by this study will contribute to a more complete picture of flavivirus biology and evolution.

Whether or not any of these viruses impact the biology of An. gambiae or other mosquitoes remains to be determined, as does their potential utility as transmission reducing agents. The microbiome of mosquito species has been demonstrated to alter vector competence (Cook and McGraw, 2010; Suh et al., 2009; Weiss and Aksoy, 2011; Zabalou et al., 2004). But the effect of ISVs on this is not as well studied and published results are mixed. Kenney et al. demonstrated that co-infection by Nhumirim virus, a dual host associated flavivirus, markedly reduced the production of West Nile virus and other pathogenic flaviviruses in cell culture (Kenney et al., 2014). However, other studies measuring the impact of ISV co-infection have had less promising results (Bolling et al., 2012; Kent et al., 2010; Newman et al., 2011). The identification of these anopheles viruses will enable researchers to begin to assess their impact on the transmission of P. falciparum parasites and other anopheles-vectored human pathogens. Isolation of these viruses in cell culture and will be an important first step in this effort.

Although we found a number of viruses that appear to infect An. gambiae, there is little doubt that this represents only a small fraction of such viruses. In fact, Carissimo and co-authors identified a cypovirus and a dicistrovirus in colony Anopheles mosquitoes and in wild mosquitoes in Senegal and Cambodia (Carissimo et al., 2016). We mainly sampled mosquito midguts and midgut contents from adult mosquitoes from West Africa, and it is possible that sampling of other tissues will reveal viruses with different tropisms. And, although in theory sequencing RNA would detect transcripts from DNA viruses it is notable that we only identified sequences of RNA viruses. It is likely that sequencing DNA-derived libraries would improve detection of DNA virus sequences. Additional sampling of different mosquito tissues from various developmental stages, and from different geographic locations, as well as utilizing alternative sequencing methods will continue to increase our understanding of the An. gambiae virome, and the impact of commensal viruses on mosquito biology.

Funding information

This project was supported in part by the CSU Infectious Disease Supercluster Grant "Xenosurveillance: A novel approach for interrogating the human-pathogen landscape in sub-Saharan Africa" awarded to DEB, GDE, BDF, and MDS. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Acknowledgments

The authors would like to thank the people of Liberia, Senegal, and Burkina-Faso for allowing this work to take place. We thank Erin Petrilli and Phil Buxton and the CSU NGS Facility for assistance with sequencing instrument operation. Tyson Claffey, Tyler Eike, Joe Strecker, Martin Yates, and Charles Heppner for IT support, and Claudia Rueckert, Alex Gendernalik, Jonathan Seaman, and Jasmine Donkoh for lab support.

Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at http://dx.doi.org/10.1016/j.virol.2016.07.031.

References

- Alout, H., Krajacich, B.J., Meyers, J.I., Grubaugh, N.D., Brackney, D.E., Kobylinski, K.C., Diclaro 2nd, J.W., Bolay, F.K., Fakoli, L.S., Diabate, A., Dabire, R.K., Bougma, R.W., Foy, B.D., 2014. Evaluation of ivermectin mass drug administration for malaria transmission control across different West African environments. Malar. J. 13, 417.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. J. Mol. Biol. 215, 403–410.
- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D. J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res. 25, 3389–3402.
- Attoui, H., Mohd Jaafar, F., Belhouchet, M., Biagini, P., Cantaloube, J.-F., de Micco, P., de Lamballerie, X., 2005. Expansion of family Reoviridae to include nine-segmented dsRNA viruses: Isolation and characterization of a new virus designated aedes pseudoscutellaris reovirus assigned to a proposed genus (Dinovernavirus). Virology 343, 212–223.
- Auguste, A.J., Carrington, C.V., Forrester, N.L., Popov, V.L., Guzman, H., Widen, S.G., Wood, T.G., Weaver, S.C., Tesh, R.B., 2014. Characterization of a novel Negevirus and a novel Bunyavirus isolated from Culex (Culex) declarator mosquitoes in Trinidad. J. Gen. Virol. 95, 481–485.
- Auguste, A.J., Kaelber, J.T., Fokam, E.B., Guzman, H., Carrington, C.V., Erasmus, J.H., Kamgang, B., Popov, V.L., Jakana, J., Liu, X., Wood, T.G., Widen, S.G., Vasilakis, N., Tesh, R.B., Chiu, W., Weaver, S.C., 2015. A newly isolated reovirus has the simplest genomic and structural organization of any reovirus. J. Virol. 89, 676–687.
- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N., Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J. Comput. Biol.: J. Comput. Mol. Cell Biol. 19, 455–477.
- Blitvich, B.J., Firth, A.E., 2015. Insect-specific flaviviruses: a systematic review of their discovery, host range, mode of transmission, superinfection exclusion potential and genomic organization. Viruses 7, 1927–1959.
- Boissière, A., Tchioffo, M.T., Bachar, D., Abate, L., Marie, A., Nsango, S.E., Shahbazkia, H.R., Awono-Ambene, P.H., Levashina, E.A., Christen, R., Morlais, I., 2012. Midgut microbiota of the malaria mosquito vector < italic > Anopheles gambiae </ italic > and interactions with < italic > Plasmodium falciparum </italic > Infection. PLoS Pathog. 8, e1002742.
- Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinforma. (Oxf., Engl.) 30, 2114–2120.
- Bolling, B., Weaver, S., Tesh, R., Vasilakis, N., 2015. Insect-specific virus discovery: significance for the arbovirus community. Viruses 7, 2851.
- Bolling, B.G., Olea-Popelka, F.J., Eisen, L., Moore, C.G., Blair, C.D., 2012. Transmission dynamics of an insect-specific flavivirus in a naturally infected Culex pipiens laboratory colony and effects of co-infection on vector competence for West Nile virus. Virology 427, 90–97.
- Buck, M., Nilsson, L.K., Brunius, C., Dabire, R.K., Hopkins, R., Terenius, O., 2016. Bacterial associations reveal spatial population dynamics in Anopheles gambiae mosquitoes. Sci. Rep. 6, 22806.
- Carissimo, G., Eiglmeier, K., Reveillaud, J., Holm, I., Diallo, M., Diallo, D., Vantaux, A., Kim, S., Ménard, D., Siv, S., Belda, E., Bischoff, E., Antoniewski, C., Vernick, K.D., 2016. Identification and characterization of two novel RNA viruses from anopheles gambiae species complex mosquitoes. PloS One 11, e0153881.
- Chandler, J.A., Liu, R.M., Bennett, S.N., 2015. RNA shotgun metagenomic sequencing of northern California (USA) mosquitoes uncovers viruses, bacteria, and fungi. Front. Microbiol. 6, 185.
- Chandler, J.A., Thongsripong, P., Green, A., Kittayapong, P., Wilcox, B.A., Schroth, G.P., Kapan, D.D., Bennett, S.N., 2014. Metagenomic shotgun sequencing of a Bunyavirus in wild-caught Aedes aegypti from Thailand informs the evolutionary and genomic history of the Phleboviruses. Virology 464–465, 312–319.
- Cirimotich, C.M., Dong, Y., Clayton, A.M., Sandiford, S.L., Souza-Neto, J.A., Mulenga, M., Dimopoulos, G., 2011. Natural microbe-mediated refractoriness to Plasmodium infection in Anopheles gambiae. Science 332, 855–858.
- Cook, P.E., McGraw, E.A., 2010. Wolbachia pipientis: an expanding bag of tricks to explore for disease control. Trends Parasitol. 26, 373–375.
- Cook, S., Bennett, S.N., Holmes, E.C., De Chesse, R., Moureau, G., de Lamballerie, X., 2006. Isolation of a new strain of the flavivirus cell fusing agent virus in a natural mosquito population from Puerto Rico. J. Gen. Virol. 87, 735–748.
- Cook, S., Chung, B.Y., Bass, D., Moureau, G., Tang, S., McAlister, E., Culverwell, C.L., Glucksman, E., Wang, H., Brown, T.D., Gould, E.A., Harbach, R.E., de Lamballerie, X., Firth, A.E., 2013. Novel virus discovery and genome reconstruction from field RNA samples reveals highly divergent viruses in dipteran hosts. PloS One 8, e80720.
- Cook, S., Moureau, G., Kitchen, A., Gould, E.A., de Lamballerie, X., Holmes, E.C., Harbach, R.E., 2012. Molecular evolution of the insect-specific flaviviruses. J. Gen. Virol. 93, 223–234.
- Crochu, S., Cook, S., Attoui, H., Charrel, R.N., De Chesse, R., Belhouchet, M., Lemasson, J.J., de Micco, P., de Lamballerie, X., 2004. Sequences of flavivirus-related RNA viruses persist in DNA form integrated in the genome of Aedes spp. mosquitoes. J. Gen. Virol. 85, 1971–1980.
- Dantas, M.D., Cavalcante, G.H., Oliveira, R.A., Lanza, D.C., 2016. New insights about ORF1 coding regions support the proposition of a new genus comprising arthropod viruses in the family Totiviridae. Virus Res. 211, 159–164.
- Dong, Y., Manfredini, F., Dimopoulos, G., 2009. Implication of the mosquito midgut

microbiota in the defense against malaria parasites. PLoS Pathog. 5, e1000423. Fauver, J.R., Pecher, L., Schurich, J.A., Bolling, B.G., Calhoon, M., Grubaugh, N.D.,

- Burkhalter, K.L., Eisen, L., Andre, B.G., Nasci, R.S., LeBailly, A., Ebel, G.D., Moore, C.G., 2015. Temporal and spatial variability of entomological risk indices for west nile virus infection in Northern Colorado: 2006–2013. J. Med. Entomol. Net ACC Distribution of the UNIVERSITY of the Colorado Colorado
- Firth, A.E., Blitvich, B.J., Wills, N.M., Miller, C.L., Atkins, J.F., 2010. Evidence for ribosomal frameshifting and a novel overlapping gene in the genomes of insectspecific flaviviruses. Virology 399, 153–166.
- Gendrin, M., Rodgers, F.H., Yerbanga, R.S., Ouedraogo, J.B., Basanez, M.G., Cohuet, A., Christophides, G.K., 2015. Antibiotics in ingested human blood affect the mosquito microbiota and capacity to transmit malaria. Nat. Commun. 6, 5921.
- Grubaugh, N.D., Massey, A., Shives, K.D., Stenglein, M.D., Ebel, G.D., Beckham, J.D., 2016. West nile virus population structure, injury, and interferon-stimulated gene expression in the brain from a fatal case of encephalitis. Open Forum Infect. Dis. 3, ofv182.
- Grubaugh, N.D., Sharma, S., Krajacich, B.J., Fakoli Iii, L.S., Bolay, F.K., Diclaro Ii, J.W., Johnson, W.E., Ebel, G.D., Foy, B.D., Brackney, D.E., 2015. Xenosurveillance: a novel mosquito-based approach for examining the human-pathogen landscape. PLoS Negl, Trop. Dis. 9, e0003628.
- Hermanns, K., Zirkel, F., Kurth, A., Drosten, C., Junglen, S., 2014. Cimodo virus belongs to a novel lineage of reoviruses isolated from African mosquitoes. J. Gen. Virol. 95, 905–909.
- Holt, R.A., Subramanian, G.M., Halpern, A., Sutton, G.G., Charlab, R., Nusskern, D.R., Wincker, P., Clark, A.G., Ribeiro, J.M., Wides, R., Salzberg, S.L., Loftus, B., Yandell, M., Majoros, W.H., Rusch, D.B., Lai, Z., Kraft, C.L., Abril, J.F., Anthouard, V., Are-nsburger, P., Atkinson, P.W., Baden, H., de Berardinis, V., Baldwin, D., Benes, V., Biedler, J., Blass, C., Bolanos, R., Boscus, D., Barnstead, M., Cai, S., Center, A., Chaturverdi, K., Christophides, G.K., Chrystal, M.A., Clamp, M., Cravchik, A., Curwen, V., Dana, A., Delcher, A., Dew, I., Evans, C.A., Flanigan, M., Grund-schober-Freimoser, A., Friedli, L., Gu, Z., Guan, P., Guigo, R., Hillenmeyer, M.E., Hladun, S.L., Hogan, J.R., Hong, Y.S., Hoover, J., Jaillon, O., Ke, Z., Kodira, C., Kokoza, E., Koutsos, A., Letunic, I., Levitsky, A., Liang, Y., Lin, J.J., Lobo, N.F., Lopez, J. R., Malek, J.A., McIntosh, T.C., Meister, S., Miller, J., Mobarry, C., Mongin, E., Murphy, S.D., O'Brochta, D.A., Pfannkoch, C., Qi, R., Regier, M.A., Remington, K., Shao, H., Sharakhova, M.V., Sitter, C.D., Shetty, J., Smith, T.J., Strong, R., Sun, J., Thomasova, D., Ton, L.Q., Topalis, P., Tu, Z., Unger, M.F., Walenz, B., Wang, A., Wang, J., Wang, M., Wang, X., Woodford, K.J., Wortman, J.R., Wu, M., Yao, A., Zdobnov, E.M., Zhang, H., Zhao, Q., Zhao, S., Zhu, S.C., Zhimulev, I., Coluzzi, M., della Torre, A., Roth, C.W., Louis, C., Kalush, F., Mural, R.J., Myers, E.W., Adams, M.D., Smith, H.O., Broder, S., Gardner, M.J., Fraser, C.M., Birney, E., Bork, P., Brey, P.T., Venter, J.C., Weissenbach, J., Kafatos, F.C., Collins, F.H., Hoffman, S.L., 2002. The genome sequence of the malaria mosquito Anopheles gambiae. Science 298, 129–149.
- Hoshino, K., Isawa, H., Tsuda, Y., Yano, K., Sasaki, T., Yuda, M., Takasaki, T., Kobayashi, M., Sawabe, K., 2007. Genetic characterization of a new insect flavivirus isolated from Culex pipiens mosquito in Japan. Virology 359, 405–414.
- Huang, Y., Mi, Z., Zhuang, L., Ma, M., An, X., Liu, W., Cao, W., Tong, Y., 2013. Presence of entomobirnaviruses in Chinese mosquitoes in the absence of Dengue virus co-infection. J. Gen. Virol. 94, 663–667.
- Hughes, G.L., Rivero, A., Rasgon, J.L., 2014. Wolbachia can enhance plasmodium infection in mosquitoes: implications for malaria control? PLoS Pathog. 10, e1004182.
- Isawa, H., Kuwata, R., Hoshino, K., Tsuda, Y., Sakai, K., Watanabe, S., Nishimura, M., Satho, T., Kataoka, M., Nagata, N., Hasegawa, H., Bando, H., Yano, K., Sasaki, T., Kobayashi, M., Mizutani, T., Sawabe, K., 2011. Identification and molecular characterization of a new non segmented double-stranded RNA virus isolated from Culex mosquitoes in Japan. Virus Res. 155, 147–155.
- Junglen, S., Drosten, C., 2013. Virus discovery and recent insights into virus diversity in arthropods. Curr. Opin. Microbiol. 16, 507–513.
- Kallies, R., Kopp, A., Zirkel, F., Estrada, A., Gillespie, T.R., Drosten, C., Junglen, S., 2014. Genetic characterization of goutanap virus, a novel virus related to negeviruses, cileviruses and higreviruses. Viruses 6, 4346–4357.
- Katoh, K., Standley, D.M., 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Mol. Biol. Evol. 30, 772–780.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond, A., 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformation 28, 1647–1649.
- Kenney, J.L., Solberg, O.D., Langevin, S.A., Brault, A.C., 2014. Characterization of a novel insect-specific flavivirus from Brazil: potential for inhibition of infection of arthropod cells with medically important flaviviruses. J. Gen. Virol. 95, 2796–2808.
- Kent, R.J., Crabtree, M.B., Miller, B.R., 2010. Transmission of West Nile virus by Culex quinquefasciatus say infected with Culex Flavivirus Izabal. PLoS Negl. Trop. Dis. 4, e671.
- Koyama, S., Urayama, S., Ohmatsu, T., Sassa, Y., Sakai, C., Takata, M., Hayashi, S., Nagai, M., Furuya, T., Moriyama, H., Satoh, T., Ono, S., Mizutani, T., 2015. Identification, characterization and full-length sequence analysis of a novel dsRNA virus isolated from the arboreal ant Camponotus yamaokai. J. Gen. Virol. 96, 1930–1937.
- Krajacich, B.J., Slade, J.R., Mulligan, R.F., LaBrecque, B., Alout, H., Grubaugh, N.D., Meyers, J.I., Fakoli 3rd, L.S., Bolay, F.K., Brackney, D.E., Burton, T.A., Seaman, J.A., Diclaro 2nd, J.W., Dabire, R.K., Foy, B.D., 2015. Sampling host-seeking anthropophilic mosquito vectors in west Africa: comparisons of an active human-

baited tent-trap against gold standard methods. Am. J. Trop. Med. Hyg. 92, 415–421.

- Krishna, S.S., Majumdar, I., Grishin, N.V., 2003. Structural classification of zinc fingers: survey and summary. Nucleic Acids Res. 31, 532–550.
- Kuhn, J.H., Bekal, S., Caì, Y., Clawson, A.N., Domier, L.L., Herrel, M., Jahrling, P.B., Kondo, H., Lambert, K.N., Mihindukulasuriya, K.A., Nowotny, N., Radoshitzky, S. R., Schneider, U., Staeheli, P., Suzuki, N., Tesh, R.B., Wang, D., Wang, L.-F., Dietzgen, R.G., 2013. Nyamiviridae: proposal for a new family in the order Mononegavirales. Arch. Virol. 158. http://dx.doi.org/10.1007/ s00705-00013-01674-y.
- Kuhn, J.H., Durrwald, R., Bao, Y., Briese, T., Carbone, K., Clawson, A.N., deRisi, J.L., Garten, W., Jahrling, P.B., Kolodziejek, J., Rubbenstroth, D., Schwemmle, M., Stenglein, M., Tomonaga, K., Weissenbock, H., Nowotny, N., 2015. Taxonomic reorganization of the family Bornaviridae. Arch. Virol. 160, 621–632.
- Kuno, G., Chang, G.J., Tsuchiya, K.R., Karabatsos, N., Cropp, C.B., 1998. Phylogeny of the genus Flavivirus. J. Virol. 72, 73–83.
- Kuwata, R., Isawa, H., Hoshino, K., Tsuda, Y., Yanase, T., Sasaki, T., Kobayashi, M., Sawabe, K., 2011. RNA splicing in a new rhabdovirus from Culex mosquitoes. J. Virol. 85, 6185–6196.
- Kuwata, R., Satho, T., Isawa, H., Yen, N.T., Phong, T.V., Nga, P.T., Kurashige, T., Hiramatsu, Y., Fukumitsu, Y., Hoshino, K., Sasaki, T., Kobayashi, M., Mizutani, T., Sawabe, K., 2013. Characterization of Dak Nong virus, an insect nidovirus isolated from Culex mosquitoes in Vietnam. Arch. Virol. 158, 2273–2284.
- Ladner, J.T., Beitzel, B., Chain, P.S., Davenport, M.G., Donaldson, E.F., Frieman, M., Kugelman, J.R., Kuhn, J.H., O'Rear, J., Sabeti, P.C., Wentworth, D.E., Wiley, M.R., Yu, G.Y., Sozhamannan, S., Bradburne, C., Palacios, G., 2014. Standards for sequencing viral genomes in the era of high-throughput sequencing. mBio 5, e01360–01314.
- Lanciotti, R.S., Ludwig, M.L., Rwaguma, E.B., Lutwama, J.J., Kram, T.M., Karabatsos, N., Cropp, B.C., Miller, B.R., 1998. Emergence of Epidemic O'nyong-nyong fever in Uganda after a 35-Year absence: genetic characterization of the virus. Virology 252, 258–268.
- Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods 9, 357–359.
- Lee, J.S., Grubaugh, N.D., Kondig, J.P., Turell, M.J., Kim, H.C., Klein, T.A., O'Guinn, M.L., 2013. Isolation and genomic characterization of Chaoyang virus strain ROK144 from Aedes vexans nipponii from the Republic of Korea. Virology 435, 220–224.
- Li, C.X., Shi, M., Tian, J.H., Lin, X.D., Kang, Y.J., Chen, L.J., Qin, X.C., Xu, J., Holmes, E.C., Zhang, Y.Z., 2015. Unprecedented genomic diversity of RNA viruses in arthropods reveals the ancestry of negative-sense RNA viruses. eLife, 4.
- Li, W., Godzik, A., 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformation 22, 1658–1659.
- Lutwama, J.J., Kayondo, J., Savage, H.M., Burkot, T.R., Miller, B.R., 1999. Epidemic O'Nyong-Nyong fever in southcentral Uganda, 1996–1997: entomologic studies in Bbaale village, Rakai District. Am. J. Trop. Med. Hyg. 61, 158–162.
- Marklewitz, M., Gloza-Rausch, F., Kurth, A., Kummerer, B.M., Drosten, C., Junglen, S., 2012. First isolation of an Entomobirnavirus from free-living insects. J. Gen. Virol. 93, 2431–2435.
- Marklewitz, M., Handrick, S., Grasse, W., Kurth, A., Lukashev, A., Drosten, C., Ellerbrok, H., Leendertz, F.H., Pauli, G., Junglen, S., 2011. Gouleako virus isolated from West African mosquitoes constitutes a proposed novel genus in the family Bunyaviridae. J. Virol. 85, 9227–9234.
- Marklewitz, M., Zirkel, F., Rwego, I.B., Heidemann, H., Trippner, P., Kurth, A., Kallies, R., Briese, T., Lipkin, W.I., Drosten, C., Gillespie, T.R., Junglen, S., 2013. Discovery of a unique novel clade of mosquito-associated bunyaviruses. J. Virol. 87, 12850–12865.
- Martinez, J., Lepetit, D., Ravallec, M., Fleury, F., Varaldi, J., 2016. Additional heritable virus in the parasitic wasp Leptopilina boulardi: prevalence, transmission and phenotypic effects. J. Gen. Virol. 97, 523–535.
- Mihindukulasuriya, K.A., Nguyen, N.L., Wu, G., Huang, H.V., da Rosa, A.P., Popov, V.L., Tesh, R.B., Wang, D., 2009. Nyamanini and midway viruses define a novel taxon of RNA viruses in the order Mononegavirales. J. Virol. 83, 5109–5116.
- Misencik, M.J., Grubaugh, N.D., Andreadis, T.G., Ebel, G.D., Armstrong, P.M., 2016. Isolation of a novel insect-specific flavivirus from culiseta melanura in the Northeastern United States. Vector Borne Zoonotic Dis. 16, 181–190.
- Nabeshima, T., Inoue, S., Okamoto, K., Posadas-Herrera, G., Yu, F., Uchida, L., Ichinose, A., Sakaguchi, M., Sunahara, T., Buerano, C.C., Tadena, F.P., Orbita, I.B., Natividad, F.F., Morita, K., 2014. Tanay virus, a new species of virus isolated from mosquitoes in the Philippines. J. Gen. Virol. 95, 1390–1395.
- Nasar, F., Palacios, G., Gorchakov, R.V., Guzman, H., Da Rosa, A.P., Savji, N., Popov, V. L., Sherman, M.B., Lipkin, W.I., Tesh, R.B., Weaver, S.C., 2012. Eilat virus, a unique alphavirus with host range restricted to insects by RNA replication. Proc. Natl. Acad. Sci. USA 109, 14622–14627.
- Newman, C.M., Cerutti, F., Anderson, T.K., Hamer, G.L., Walker, E.D., Kitron, U.D., Ruiz, M.O., Brawn, J.D., Goldberg, T.L., 2011. Culex flavivirus and West Nile virus mosquito co infection and positive ecological association in Chicago. U. S. Vector Borne Zoonotic Dis. 11, 1099–1105.
- Nga, P.T., Parquet, Md.C., Lauber, C., Parida, M., Nabeshima, T., Yu, F., Thuy, N.T., Inoue, S., Ito, T., Okamoto, K., Ichinose, A., Snijder, E.J., Morita, K., Gorbalenya, A. E., 2011. Discovery of the first insect nidovirus, a missing evolutionary link in the emergence of the largest RNA virus genomes. PLoS Pathog. 7, e1002215.
- Poulos, B.T., Tang, K.F., Pantoja, C.R., Bonami, J.R., Lightner, D.V., 2006. Purification and characterization of infectious myonecrosis virus of penaeid shrimp. J. Gen. Virol. 87, 987–996.
- Pybus, O.G., Rambaut, A., Holmes, E.C., Harvey, P.H., 2002. New inferences from tree shape: numbers of missing taxa and population growth rates. Syst. Biol. 51,

881-888.

- Quan, P.L., Junglen, S., Tashmukhamedova, A., Conlan, S., Hutchison, S.K., Kurth, A., Ellerbrok, H., Egholm, M., Briese, T., Leendertz, F.H., Lipkin, W.I., 2010. Moussa virus: a new member of the Rhabdoviridae family isolated from Culex decens mosquitoes in Cote d'Ivoire. Virus Res. 147, 17–24.
- Ren, X., Hoiczyk, E., Rasgon, J.L., 2008. Viral paratransgenesis in the malaria vector Anopheles gambiae. PLoS Pathog. 4, e1000135.
- Ronquist, F., Huelsenbeck, J.P., 2003. MrBayes 3: bayesian phylogenetic inference under mixed models. Bioinformation 19, 1572–1574.
- Schuster, S., Zirkel, F., Kurth, A., van Cleef, K.W., Drosten, C., van Rij, R.P., Junglen, S., 2014. A unique nodavirus with novel features: mosinovirus expresses two subgenomic RNAs, a capsid gene of unknown origin, and a suppressor of the antiviral RNA interference pathway. J. Virol. 88, 13447–13459.
- Soding, J., Biegert, A., Lupas, A.N., 2005. The HHpred interactive server for protein homology detection and structure prediction. Nucleic Acids Res. 33, W244–W248.
- Spear, A., Sisterson, M.S., Yokomi, R., Stenger, D.C., 2010. Plant-feeding insects harbor double-stranded RNA viruses encoding a novel proline-alanine rich protein and a polymerase distantly related to that of fungal viruses. Virology 404, 304–311.
- Suh, E., Mercer, D.R., Fu, Y., Dobson, S.L., 2009. Pathogenicity of life-shortening Wolbachia in Aedes albopictus after transfer from Drosophila melanogaster. Appl. Environ. Microbiol. 75, 7783–7788.
- Suzuki, Y., Niu, G., Hughes, G.L., Rasgon, J.L., 2014. A viral over-expression system for the major malaria mosquito Anopheles gambiae. Sci. Rep. 4, 5127.
- Talavera, G., Castresana, J., 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. Syst. Biol. 56, 564–577.
- Tokarz, R., Williams, S.H., Sameroff, S., Sanchez Leon, M., Jain, K., Lipkin, W.I., 2014. Virome analysis of Amblyomma americanum, Dermacentor variabilis, and lxodes scapularis ticks reveals novel highly divergent vertebrate and invertebrate viruses. J. Virol. 88, 11480–11492.
- Tyler, S., Bol^Ting, B.G., Blair, C.D., Brault, A.C., Pabbaraju, K., Armijos, M.V., Clark, D.C., Calisher, C.H., Drebot, M.A., 2011. Distribution and phylogenetic comparisons of a novel mosquito flavivirus sequence present in Culex tarsalis Mosquitoes from western Canada with viruses isolated in California and Colorado. Am. J. Trop. Med. Hyg. 85, 162–168.
- Van Regenmortel, M.H.V., Fauquet, C.M., Bishop, D.H.L., Carstens, E.B., Estes, M.K., Lemon, S.M., Maniloff, J., Mayo, M.A., McGeoch, D.J., Pringle, C.R., Wickner, R.B., 2000a. Order Mononegavirales, Virus Taxonomy. Seventh Report of the International Committee on Taxonomy of Viruses Academic Press, pp. 525–530.
- Van Regenmortel, M.H.V., Fauquet, C.M., Bishop, D.H.L., Carstens, E.B., Estes, M.K., Lemon, S.M., Maniloff, J., Mayo, M.A., McGeoch, D.J., Pringle, C.R., Wickner, R.B., 2000b. Order Mononegavirales, Virus Taxonomy. Seventh Report of the International Committee on Taxonomy of Viruses Academic Press, pp. 525–530.
- Vancini, R., Paredes, A., Ribeiro, M., Blackburn, K., Ferreira, D., Kononchik Jr., J.P., Hernandez, R., Brown, D., 2012. Espirito Santo virus: a new birnavirus that replicates in insect cells. J. Virol. 86, 2390–2399.
- Vasilakis, N., Castro-Llanos, F., Widen, S.G., Aguilar, P.V., Guzman, H., Guevara, C., Fernandez, R., Auguste, A.J., Wood, T.G., Popov, V., Mundal, K., Ghedin, E., Kochel, T.J., Holmes, E.C., Walker, P.J., Tesh, R.B., 2014a. Arboretum and Puerto Almendras viruses: two novel rhabdoviruses isolated from mosquitoes in Peru. J. Gen. Virol. 95, 787–792.

- Vasilakis, N., Forrester, N.L., Palacios, G., Nasar, F., Savji, N., Rossi, S.L., Guzman, H., Wood, T.G., Popov, V., Gorchakov, R., González, A.V., Haddow, A.D., Watts, D.M., da Rosa, A.P.A.T., Weaver, S.C., Lipkin, W.I., Tesh, R.B., 2013. Negevirus: a proposed new taxon of insect-specific viruses with wide geographic distribution. J. Virol. 87, 2475–2488.
- Vasilakis, N., Guzman, H., Firth, C., Forrester, N.L., Widen, S.G., Wood, T.G., Rossi, S.L., Ghedin, E., Popov, V., Blasdell, K.R., Walker, P.J., Tesh, R.B., 2014b. Mesoniviruses are mosquito-specific viruses with extensive geographic distribution and host range. Virol. J. 11, 97.
- Vasilakis, N., Tesh, R.B., 2015. Insect-specific viruses and their potential impact on arbovirus transmission. Curr. Opin. Virol. 15, 69–74.
- Wang, L., Lv, X., Zhai, Y., Fu, S., Wang, D., Rayner, S., Tang, Q., Liang, G., 2012. Genomic characterization of a novel virus of the family <italic > Tymoviridae </italic > isolated from mosquitoes. PloS One 7, e39845.
- Warrilow, D., Watterson, D., Hall, R.A., Davis, S.S., Weir, R., Kurucz, N., Whelan, P., Allcock, R., Hall-Mendelin, S., O'Brien, C.A., Hobson-Peters, J., 2014. A new species of mesonivirus from the Northern Territory, Australia. PloS One 9, e91103.
- Weiss, B., Aksoy, S., 2011. Microbiome influences on insect host vector competence. Trends Parasitol. 27, 514–522.
- World Health Organization, 2015. World Malaria Report.
- World Health Organization, 2016. Lymphatic filariasis Epidemiology.
- Wu, Q., Luo, Y., Lu, R., Lau, N., Lai, E.C., Li, W.X., Ding, S.W., 2010. Virus discovery by deep sequencing and assembly of virus-derived small silencing RNAs. Proc. Natl. Acad. Sci. USA 107, 1606–1611.
- Wu, T.D., Nacu, S., 2010. Fast and SNP-tolerant detection of complex variants and splicing in short reads. Bioinformation 26, 873–881.
- Yamao, T., Eshita, Y., Kihara, Y., Satho, T., Kuroda, M., Sekizuka, T., Nishimura, M., Sakai, K., Watanabe, S., Akashi, H., Rongsriyam, Y., Komalamisra, N., Srisawat, R., Miyata, T., Sakata, A., Hosokawa, M., Nakashima, M., Kashige, N., Miake, F., Fukushi, S., Nakauchi, M., Saijo, M., Kurane, I., Morikawa, S., Mizutani, T., 2009. Novel virus discovery in field-collected mosquito larvae using an improved system for rapid determination of viral RNA sequences (RDV ver4.0). Arch. Virol. 154, 153–158.
- Zabalou, S., Riegler, M., Theodorakopoulou, M., Stauffer, C., Savakis, C., Bourtzis, K., 2004. Wolbachia-induced cytoplasmic incompatibility as a means for insect pest population control. Proc. Natl. Acad. Sci. USA 101, 15042–15045.
- Zhai, Y., Attoui, H., Mohd Jaafar, F., Wang, H.Q., Cao, Y.X., Fan, S.P., Sun, Y.X., Liu, L.D., Mertens, P.P., Meng, W.S., Wang, D., Liang, G., 2010. Isolation and full-length sequence analysis of Armigeres subalbatus totivirus, the first totivirus isolate from mosquitoes representing a proposed novel genus (Artivirus) of the family Totiviridae. J. Gen. Virol. 91, 2836–2845.
- Zhao, Y., Tang, H., Ye, Y., 2012. RAPSearch2: a fast and memory-efficient protein similarity search tool for next-generation sequencing data. Bioinformation 28, 125–126.
- Zirkel, F., Kurth, A., Quan, P.L., Briese, T., Ellerbrok, H., Pauli, G., Leendertz, F.H., Lipkin, W.I., Ziebuhr, J., Drosten, C., Junglen, S., 2011. An insect nidovirus emerging from a primary tropical rainforest. mBio 2, e00077-00011.
- Zirkel, F., Roth, H., Kurth, A., Drosten, C., Ziebuhr, J., Junglen, S., 2013. Identification and characterization of genetically divergent members of the newly established family Mesoniviridae. J. Virol. 87, 6346–6358.